Principle of Interferometer Beam Splitter



Overview

Using a beam splitter, a light source is split into two arms. The Michelson interferometer is a common configuration for optical interferometry and was invented by the American physicist Albert Abraham Michelson in 1887. Each of those light beams is reflected back toward the beamsplitter which then. erferometer employs a division of amplitude scheme. A beam splitter is nothing more than a plate of glass, which is made partially reflective: as. A Michelson interferometer, as invented by Albert Abraham Michelson in the late 19 th century, is a very common type of interferometer, which is frequently used for measuring distances or their changes, but also in various other applications (see below). While in a spectrometer (see the hydrogen spectrum.

Article Content

1. Introduction 2. Michelson interferometer: theory

he original width by having reduced amplitudes . A beam splitter is nothing more than a plate of glass, which is made partially reflective: as such, the splitting occurs because part of the light is reflected off.

Michelson Interferometers – operation principle, arm

A Michelson interferometer (see Figure 1) contains a single beam splitter for separating and recombining the beams. There are two interferometer arms (each

Michelson Interferometry

© 2026 KENHARDT OPTICAL NETWORKS - All rights reserved

The Michelson Interferometer is an amplitude-splitting interferometer. It splits the beam into two perpendicular paths using a 50 % beam splitter. The beams then reflect from the mirrors placed on

Mach-Zehnder interferometer

A Mach-Zehnder interferometer consists of two beam splitters (thick grey lines) and two mirrors (thin lines). A light beam splits at the first beam splitter, travels some distance in top top and

16. Michelson Interferometer — Modern Lab

Light from a laser is incident on a beam splitter (BS) which consists of a glass plate with a partially reflective surface. About 50% of the light is reflected

Infrared Spectroscopy: Beam Splitters and Detector Physics Explained

A broadband infrared source hits a beam splitter, which splits the light into two paths—one heads to a fixed mirror, the other to a moving mirror. The reflected beams meet up again at the

Interferometers 1. Introduction 2. Michelson interferometer: theory

Differences in the velocity of light along 2 different directions. It operates as follows: we “divide” the wave amplitude by partial reflection using a beam splitter G1, with the two resulting wave fro.

Interferometers

According to this principle, the incident beam of light falls on a beam splitter, which reflects roughly half of the intensity of the wave front in one direction and transmits the other half of the intensity of the

The Michelson Interferometer

These instruments are grouped under the generic name of interferometers. The Michelson interferometer causes interference by splitting a beam of light into two parts. Each part is made to

Understanding Beamsplitters: Types, Principles, and

This article explores the fundamental principles and diverse applications of beamsplitters, detailing their different types and uses in fields

1. Introduction 2. Michelson interferometer: theory

The two beams obtained by amplitude division are sent in different directions against plane mirrors, then reflected back along their same respected paths to the beam splitter to form an interference pattern.

© 2026 KENHARDT OPTICAL NETWORKS - All rights reserved

Beamsplitter

To calculate the intensities of the combined beams falling on the detector and on the source, we start by considering the phase difference between the reflected and transmitted beams leaving the

Michelson Interferometer: Ultimate Guide

Explore the world of optical instrumentation with our comprehensive guide to Michelson Interferometer, covering its principles, applications, and significance.

How Beamsplitters Work: Types, Mechanisms, and

This article explains the working principles of beamsplitters, detailing how they divide a beam of light into two separate paths, the different types of

Chapter 4: The Michelson Interferometer

4.3 Working Principle When light from the source reaches the beam splitter, it splits into two beams traveling along different paths, reflected by the mirrors. The beams then recombine at the

Beamsplitter

Beamsplitter The beamsplitter is one of the most expensive and sensitive components of an interferometer, and must be chosen carefully. A pellicle beamsplitter is a high tensile strength elastic

How do interferometers work?

How do interferometers work? An interferometer is a really precise scientific instrument designed to measure things with extraordinary accuracy.

Interferometry

Fabry-Perot interferometer: theory7 Another commonplace division-of-amplitude interferometer is the Fabry-Perot inter-ferometer, which uses a principle similar to that of the Michelson interferometer to

Related Video Reference

This video was associated with the source search result. Verify technical details against current product documentation and project requirements.

© 2026 KENHARDT OPTICAL NETWORKS - All rights reserved

Contact Us

Continue your research online:

Website: https://kenhardtprimary.co.za
Technical inquiry: https://kenhardtprimary.co.za/project-desk.html
African project coordination: Century City, Cape Town, Western Cape, South Africa

This document is for preliminary research. Verify specifications against current standards, product documentation and project requirements.

© 2026 KENHARDT OPTICAL NETWORKS - All rights reserved